## **NEW PERIODONTAL CLASSIFICATION QUICK REFERENCE GUIDE**



The following 5 steps must be followed to create a patient 'Periodontal Diagnostic Statement'

TYPE: STEP 1 Periodontally healthy, gingivitis or periodontitis? **EXTENT:** What is the pattern of the disease, how many and what teeth are affected? **CURRENT DISEASE STATE:** STEP 3 Based on probing depths and BoP STAGING: STEP 4 This tells us how severe the disease is. Look at radiographs for the site with the worst interproximal bone loss **GRADING:** STEP 5 This is about disease susceptibility and tells us how fast the disease is progressing. Look at radiographs for the site with the worst interproximal bone loss and divide the % bone loss by

Periodontal health – no obvious evidence of interdental recession

**⇔ Gingivitis** – no obvious evidence of interdental recession

**→ Periodontitis** – progress to step 2

**Generalised** – affects ≥ 30% of teeth

► Localised – <30% of teeth

■ Molar incisal pattern

Stable – BoP <10%, PPD ≤4mm, No BoP at 4mm sites

•• In remission – BoP ≥ 10%, PPD ≤4mm, No BoP at 4mm sites

Unstable - PPD ≥5mm or PPD at ≥4mm & BoP

■ Stage I – <15% (or <2mm attachment loss from CEJ)

■ Stage II – coronal third of root

**Stage III** − mid third of root

**Stage IV** − apical third of root

**Grade A (slow)** − < 0.5

**⇔ Grade B (moderate)** − 0.5-1.0

**⇔ Grade C (rapid)** − >1.0

TYPE: EXTENT: DISEASE STATE: STAGING: GRADING

the patient's age

## **Product Information**

Corsodyl 0.2% Mouthwash (Alcohol Free). Active Ingredient: Chlorhexidine digluconate Indications: Plaque inhibition; gingivitis; maintenance of oral hygiene; post periodontal surgery or treatment; aphthous ulceration; oral candida. Legal category: GSL. Licence Holder: GlaxoSmithKline Consumer Healthcare, Brentford, TW8 9GS, U.K. Information about this product, including adverse reactions, precautions, contraindications and method of use can be found at: https://www.medicines.org.uk/emc/medicine/23034