

100 TABLETS
(50 PACKETS OF 2 COATED TABLETS EACH)

- Headache
- Muscular Aches
- Minor Pain of Arthritis
- Menstrual Cramps
- Toothache
- Fever



Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Advil

100 TABLETS (50 PACKETS OF 2 COATED TABLETS EACH)

- Headache
- Minor Pain of Arthritis
- Muscular Aches
- Menstrual Cramps
- Fever

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Advil

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Advil

Drug Facts Active ingredient (in each tablet) Ibuprofen 200 mg (NSAID)*Pain reliever/Fever reducer *nonsteroidal anti-inflammatory drug	Drug Facts (continued) When using this product ■ take with food or milk if stomach upset occurs Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear If pregnant or breast-feeding , ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever	Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor
Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.	Other information ■ read all warnings and directions before use. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F) Inactive ingredients acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax
Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug	Questions or comments? call toll free 1-800-88-ADVIL

Do Not Use if the sealed child resistant pouch is broken or torn.

Distributed by: Haleon, Warren, NJ 07059
For most recent product information, visit www.Advil.com
Trademarks are owned by or licensed to the Haleon group of companies.
©2024 Haleon group of companies or its licensor. Patents: www.productpats.com

215102

100 TABLETS
(50 PACKETS OF 2 COATED TABLETS EACH)

- Headache
- Muscular Aches
- Minor Pain of Arthritis
- Menstrual Cramps
- Toothache
- Fever



Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Advil