

Personalized Pain Management Solutions

Our trusted OTC brands, built on science



	Advil Dual Action	Advil Liqui-Gels	Advil Coated Tablets	Excedrin Extra Strength	Excedrin Migraine
KEY FEATURES	Effective pain relief with a lower maximum daily dose of each medication	Rapid pain relief starting in minutes Available with an osteoarthritis cap	Trusted pain relief for 35+ years Available with an osteoarthritis cap	Fast-acting formula – for some, headache relief starts in as little as 15 minutes	Proven to provide effective moderate to severe migraine relief with just 1 dose Starts relieving migraine pain in 30 minutes for some patients
TOP INDICATIONS	Backache, dental pain, menstrual pain & other acute pains	Headache, musculoskeletal pain, fever, pain from common cold, & minor pain of arthritis	Minor pain of arthritis, fever, headache, pain from common cold, & musculoskeletal pain	Headaches	Migraines
INGREDIENTS (PER TABLET)	Acetaminophen 250 mg Ibuprofen 125 mg	Ibuprofen 200 mg	Ibuprofen 200 mg	Acetaminophen 250 mg Aspirin 250 mg Caffeine 65 mg	Acetaminophen 250 mg Aspirin 250 mg Caffeine 65 mg
DURATION	8 hours	6-8 hours	6-8 hours	Up to 6 hours	Up to 24 hours
DOSAGE	Adults & children 12 years & older: 2 tablets every 8 hours Maximum of 6 tablets over 24 hours	Adults and children over 12: 1 to 2 tablets every 4-6 hours as needed Maximum of 6 tablets over 24 hours	Adults and children over 12: 1 to 2 tablets every 4-6 hours as needed Maximum of 6 tablets over 24 hours	Adults and children 12 years and over: Take 2 tablets every 6 hours with a glass of water Maximum of 8 tablets over 24 hours	Adults 18+: Take 2 tablets per day with a glass of water Maximum of 2 tablets over 24 hours

For more information, including additional available products, please visit www.haleonhealthpartner.com/en-us



Keep your patients moving

Voltaren Arthritis Pain Gel

KEY FEATURES	<p>Potency Clinically proven to treat pain caused by inflammation^{1,2}</p>	<p>Penetration Deep at the source of pain^{3,4}</p>	<p>Pain-on-movement reduction^{1,5} Improves mobility^{2,3}</p>	<p>Safety profile comparable to placebo*^{1,2}</p>
	<p>Diclofenac sodium (topical NSAID) 1% indicated for relief of osteoarthritis pain in:</p>			
	<p>Hand/Wrist</p>	<p>Knee</p>	<p>Elbow</p>	<p>Foot/Ankle</p>
	<p>DURATION</p> <p>Approved for use for up to 21 days without additional intervention After 21 days, patients should consult a physician</p>			
<p>DOSING</p> <p>Patients should use Voltaren 4 times a day, every day Do not use on more than 2 body areas at the same time</p> <p>Patients should use the supplied dosing card to measure a dose:</p> <ul style="list-style-type: none"> • For each upper body area (hand, wrist, or elbow) squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle, or knee) squeeze out 4.5 inches (4 grams) 				

The American College of Rheumatology strongly recommends topical NSAIDs, like Voltaren®, as a first-line treatment of osteoarthritis pain



Scan here to learn more

*In placebo-controlled trials, adverse reactions were comparable to placebo. Application site reaction was the only AE that occurred in >1% of patients and with greater frequency for Voltaren® Arthritis Pain gel OTC vs placebo.

References: 1. Barthel HR, et al. Randomized controlled trial of diclofenac sodium gel in knee osteoarthritis. *Semin Arthritis Rheum.* 2009;39(3):203-212. 2. Altman RD, et al. Diclofenac sodium gel in patients with primary hand osteoarthritis: a randomized, double-blind, placebo-controlled trial. *J Rheumatol.* 2009;36(9):1991-1999. 3. Zacher J, et al. Topical diclofenac and its role in pain and inflammation: an evidence-based review. *Curr Med Res Opin.* 2008;24:925-950. 4. Riess W, et al. The percutaneous absorption of diclofenac. *Arzneimittelforschung.* 1986;36:1092-1096. 5. Fudin J, et al. Clinically meaningful results with a Voltaren Arthritis Pain (topical diclofenac sodium 1%), a nonsurgical treatment option for osteoarthritis of the knee. [Poster] Exhibited at PAINWeek Live Virtual Conference; September 11-13, 2020.