



Voltaren Arthritis Pain Gel is the
1st full prescription strength,
clinically-proven nonsteroidal
anti-inflammatory gel that is
approved for the treatment of
arthritis pain.



Drug Facts

Drug Facts

Active ingredient

Diclofenac sodium (NSAID*) 1% (equivalent to 0.93% diclofenac) Arthritis pain reliever
*nonsteroidal anti-inflammatory drug

Purpose

Uses

- for the temporary relief of arthritis pain **ONLY** in the following areas:
 - hand, wrist, elbow (upper body areas)
 - foot, ankle, knee (lower body areas)
- this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.

Warnings

For external use only

Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash
- If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains diclofenac. Liver damage may occur if you apply more or for a longer time than directed • when using other drugs containing diclofenac

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

Drug Facts (continued)

When using this product

- avoid contact with eyes, nose, or mouth • if eye contact occurs, rinse thoroughly with water

Stop use and ask a doctor if

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
 - feel faint • have bloody or black stools • vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain • trouble breathing • leg swelling • weakness in one part or side of body • slurred speech

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Use up to 21 days unless directed by your doctor

Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily	Per Dose
For your arthritis pain: <ul style="list-style-type: none">• Use 4 times per day every day• Do not use on more than 2 body areas at the same time	Use ENCLOSED DOSING CARD to measure a dose <ul style="list-style-type: none">• For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams)• For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)

Read the enclosed User Guide for complete instructions:

- use only as directed
- do not use more than directed or for longer than directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product
- do not apply with external heat such as heating pad • do not apply a bandage over the treated area
- store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable.

Other information

- store at 20-25°C (68-77°F). Keep from freezing.
- read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients carbomer homopolymer Type C, cocoyl caprylocaprate, fragrance, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution

Questions or comments? 1-855-297-3031 www.voltarengel.com

**Voltaren is the #1 Doctor Recommended
Topical Pain Relief Brand***

*Based on surveyed doctors, Voltaren is approved for use for the treatment of arthritis pain."

Dosing Card



Voltaren[®] ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

Upper body dose 2.25 inches long (2 grams)

Lower body dose 4.5 inches long (4 grams)

Dosing Card (NOT ACTUAL SIZE)

Read product User Guide before use
Not for immediate pain relief

The dosing card should always be used to measure a dose

- For each upper body area (hand, wrist, or elbow) squeeze out 2.25 inches (2 grams)
- For each lower body area (foot, ankle, or knee) squeeze out 4.5 inches (4 grams)

Treatment Areas & Directions

Voltaren[®] Arthritis Pain
provides effective pain relief

TREATMENT AREAS



IF YOU EXPERIENCE...

- | | | | |
|-----------------|----------------|-----------------------|---------------------------------|
| ✓
Joint pain | ✓
Stiffness | ✓
Lack of mobility | ✓
Enlarged or swollen joints |
|-----------------|----------------|-----------------------|---------------------------------|

YOU MIGHT HAVE OA

- Apply it 4 times a day, every day
- Do not use on more than 2 body areas at the same time
- After 21 days, consult a physician before further use